



Sustainable Methods for Managing Internal Parasites in Small Ruminants: Natural Dewormers

This research has enabled goat production to be a more sustainable and low-cost enterprise by mitigating internal parasites and coccidian infections of grazing animals, thus enhancing overall meat production, protecting animal and human health, increasing producer profits, and providing a stable and safe food supply.

Who cares and why?

Gastrointestinal (GI) parasites (especially *Haemonchus contortus* and *Eimeria* spp.) present the greatest danger to the goat and sheep industry in the southern United States. Infected animals have lower growth rates, reduced reproductive performance, and have higher rates of illness and death. In the past, sheep and goat producers relied heavily on anti-parasitic drugs. Unfortunately, GI parasites have become increasingly resistant to many of the ruminant anthelmintics. Alternative methods of GI parasite control for animals raised primarily on forages are vital for the sustainability and profitability of sheep and goat farms in the southern United States. Consequently, alternative, sustainable, and affordable methods of parasitic control are required. There is also a need to examine plant-based alternatives to control coccidiosis in livestock because there are no FDA approved drugs to treat coccidiosis in small ruminants.

What has the project done so far?

Research has shown that legumes such as Sericea lespedeza (*Lespedeza cuneata*) contain condensed tannin (CT) with anti-parasitic properties. The anti-parasitic properties of CT have been demonstrated to reduce GI parasitic infection in goats in Oklahoma and Georgia and in sheep in Louisiana.

Researchers at Tuskegee University found potential benefits of pine bark (PB) supplementation on anti-parasitic effects and improved feed efficiency. Pine bark is one of the abundant forest byproducts in the southern United States and contains 11-to-13 percent CT on a dry matter basis.



By raising sheep and goats primarily on forage, farmers can reduce feed costs. -Photos by Byeng R. Min, Tuskegee University.

Research demonstrates that goats fed on a PB diet have up to 30 percent fewer worms in a total ration, as well as lower instances of fecal egg count and fecal coccidian oocyst count. Feeding a pine bark diet reduced both male (64 percent) and female (59 percent) worm counts compared with the control (without PB) diet. In addition, mean dry matter intake, feed efficiency, average daily gain, and cold carcass weight were also greater for goats fed the pine bark diets.

On-farm research with local farmers in Alabama also has shown that goats on a pine bark diet had 74 percent lower fecal egg counts and 5 percent better animal weight gain compared to control diets during three-month trials. Ground pine bark as a feed ingredient has the potential to improve animal performance while decreasing internal parasites and coccidian infection. Thus, developing plant-based alternatives such as pine bark and other natural resources for GI parasites control would be expected to have a greater impact on the goat and sheep industries. This will allow development of Best Management Practices to prevent or treat coccidiosis and GI parasites in ruminant livestock.

Ultimately, by raising sheep and goats primarily on forage, farmers can reduce feed costs.

Impact Statement

Goats fed on a PB diet have up to 30 percent fewer worms in a total ration, as well as lower instances of fecal egg count and fecal coccidian oocyst count.

Goats on a pine bark diet had 74 percent lower fecal egg counts and 5 percent better animal weight gain compared to control diets during three-month trials.

Ground pine bark as a feed ingredient has the potential to improve animal performance while decreasing internal parasites and coccidian infection. Thus, developing plant-based alternatives such as pine bark and other natural resources for GI parasites control would be expected to have a greater impact on the goat and sheep industries.

Development of Best Management Practices to prevent or treat coccidiosis and GI parasites in ruminant livestock.

Raising sheep and goats primarily on forage can reduce farmers feed costs.

What research is needed?

More research is needed to: study the direct mechanisms of tannins and parasite interactions; look for ways of validating the anti-parasitic plants of sheep and goats production systems; and to continue testing the effectiveness of alternative natural dewormers for use in ruminants.

Want to know more?

Dr. Byeng Ryel Min, Research Team leader
minb@mytu.tuskegee.edu, 334-524-7670

This project was supported by the Sustainable Agriculture Research and Education (SARE) program, funded by USDA-NIFA, the USDA/NIFA Evans-Allen Research Program and Tuskegee University, George Washington Carver Agricultural Research Station.